GENERAL INDEX TO VOLUME XX

New scientific names of plants and the final members of new combinations are printed in bold face type; synonyms and page numbers having reference to figures and plates, in *italics;* and previously published scientific names and all other matter, in ordinary type.

A

Acarospora molybdina, 20; peltasticta, 20; Schleicheri, 20
Achaetogeron, 23
Acrostichum areolatum, 69
Actinea Richardsonii, 797
Adiantum, 50; Capillus-Veneris, 50; pedatum, 50
Alectoria Fremontii f. **perfertilis**, 10, subsp. olivacea, 10; implexa f. fuscidula, 10; *japonica*, 10; jubata var. prolixa, 10, var. Vrangiana, 10; laeta, 10; *lata*, 10; ochroleuca, 10; *osteina*, 10; sarmentosa, 10, var. **gigantea**, 10; *Vrangiana*, 10
Alizarin, effect of, on growth of Endomyces capsulatus, 529
Allen, Caroline K. A monograph of the American species of the genus Halenia, 119
Allomarkgrafia, 625; ovalis, 626; **plumeriaeflora**, 627
Amblyanthera, 645; *acutiloba*, 649; *andina*, 665; *Andrieuxii*, 688; *angustifolia*, 640; *antennacea*, 741; *Benthamii*, 774; *Bogotensis*, 677; *brachyloba*, 670; *Bridgesii*, 693; *campestris*, 763; *ciliata*, 763; *citrifolia*, 638; *Claussenii*, 764; *congesta*, 676; *convolvulacea*, 687; *Cuyabensis*, 752; *Fendleri*, 758; *fluminensis*, 763, α. *Stadelmeyeri*, 763, β. *Claussenii*, 763; *foliosa*, 658; *funiformis*, 697, α. *peduncularis*, 697, β. *brevipedunculata*, 697, γ. *microphylla*, 697, δ. *arenaria*, 697; *Guianensis*, 740; *hirsuta*, 763; *hispida*, 763, β. *tomentosa*, 763; *Karwinskii*, 657; *membranacea*, 744; *mexicana*, 660; *microcalyx*, 744, α. *hirtiflora*, 744, β. *trichantha*, 744, γ. *leiantha*, 744, δ. *leiophylla*, 744, ε. *glabra*, 744; *Moritziana*, 768; *Oaxacana*, 686; *ovata*, 764; *palustris*, 763, β. *almadensis*, 763; *Pavonii*, 749; *Prieurei*, 740; *Schlimi*, 759; *Spruceana*, 776; *suaveolens*, 695; *torosa*, 656; *tubiflora*, 648; *versicolor*, 752, β. *intermedia*, 755
Amesia gigantea, 802
Ammonia in relation to growth of orchid seedlings, 234, 369
Anartia Bogotensis, 677
Angadenia, 645; *Prieurii*, 740; *reticulata*, 752
Anisolobus Pohlianus, 733
Anonymus caroliniensis, 316
Aphanostephus, 23
Aplopappus gymnocephalus, 797
Apocynaceae, Studies in the. IV, 605
Aquilegia longissima, 800
Arabinose, growth of orchid seedlings in solutions containing, 574, 576
Arceuthobium vaginatum, 799
Arctostaphylos pungens, 798
Asclepias glaucescens, 796
Aspicilia calcarea, 14
Aspidium, 66; *acrostichoides*, 63, var. *incisum*, 64; *schweinitzii*, 64
Asplenium, 51; *acrostichoides*, 55; *angustifolium*, 54; *Athyrium*, 55; Bradleyi, 51; cryptolepis, 51; *ebeneum*, 53; ebenoides, 52; *Filix-foemina*, 56; *Michauxii*, 56; *parvulum*, 53; pinnatifidum, 52; platyneuron, 53, var. incisum, 53; *pycnocarpon*, 54; resiliens, 53; *Ruta-muraria*, 52; *thelypteroides*, 55; Trichomanes, 54
Astranthium, 23, 29; integrifolium, 34, *40*, var. ciliatum, 35, *44*, var. **rosulatum**, 36, *44;* **mexicanum**, 32, var. **chihuahuense**, 32, *42;* **mimum**, 30; **orthopodum**, 31; **purpurascens**, 33; **xanthocomoides**, 33; **xylopodum**, 31
Astranthium and related genera, 23
Athyrium 54; acrostichoides, 54; angustifolium, 54; angustum, 56, var. elatius, 56, var. rubellum, 56; asplenioides, 55; *Filix-foemina*, 55; *pycnocarpon*, 54; *thelypteroides*, 55
Azolla, 75; caroliniana, 75

B

Bacidia fuscorubella, 19; minuscula var. Beckhausii, 19
Bellis, 23; Garciae, 24; perennis, 23, 24, *40;* scandens, 23
Bellis, 29; *campestris*, 23; *ciliata*, 35; *integrifolia*, 34, 35, 36; *mexicana*, 31, 32; *mima*, 30; *nutans*, 34; *orthopoda*, 31; *parviflora*, 34; *pedunculata*, 23; *purpurascens*, 33; *xanthocomoides*, 33
Berkley, Dorothy Megowen and Earl E. Super optimal and thermal death temperatures of the cotton plant as affected by variations in relative humidity, 583
Biatora helvola, 19; symmictera, 19
Billardiera, 251; *explanata*, 316
Blastomyces, 103, 474; *dermatitidis*, 102, 104, 106
Blastomycoides, 85; *dermatitidis*, 104; *immitis*, 104; *tulanensis*, 104

Blastomycosis: report of case, with a study of an etiologic factor and a classification of the organism, 79; animal inoculation, 95; cultural descriptions, 98; description, 96; differential diagnosis, 90; photographs of hands affected, *118;* symptomatology, 89; technique, 95; treatment, 91
Boerhaavia erioselenus, 799
Botrychium, 47; *dissectum*, 48; obliquum, 47, var. dissectum, 48, var. tenuifolium, 48, *78; ternatum*, 47, var. *dissectum*, 48, var. *obliquum*, 47; virginianum, 48, var. intermedium, 49
Bouchea spathulata, 801
Bourdonia, 37; *bellidifolia*, 37; *effusa*, 38
Brachycome xanthocomoides, 33, 34; *zeranthemoides*, 33
Brickellia Coulteri, 797; Fendleri, 797
Bryophyta, new or rare, in flora of Texas, 795
Buchnera canadensis, 316
Buddleia marrubifolia, 799
Buellia disciformis var. insignis, 16, var. minor, 15; major, 16; punctiformis, 16

C

Calcium, relation of growth of orchid seedlings to, 232, 366
Campanula rotundifolia, 797
Camptosorus, 57; rhizophyllus, 57
Carbohydrate: reactions of Endomyces, 510; Sources of, for germination and growth of orchid seedlings, 569
Carex eburnea, 801; microdonta, 801
Catillaria tricolor, 19
Cattleya Trianae, growth of seedlings of, 224, 363, 374, 574
Ceratia, 138
Cerebrospinal meningitis, fatal, Endomyces capsulatus, a causative agent of, 471
Cetraria, 12; aleurites, 12; *californica* var. *Tuckermanii*, 11; caperata, 12; *chlorophylla*, 12; *chrysantha*, 12; cucullata, 11; glauca, 12, var. stenophylla, 12; islandica, 11; juniperina var. canadensis, 12, var. crispata, 12, var. viridis, 12; *juniperina*, var. *terrestris*, 12, var. *virescens*, 12; lacunosa var. Macounii, 12; Merrillii, 11; nigricans var. pallida, 11; nivalis, 11; norvegica, 12; *pinastri*, 12; *platyphylla*, 13; scutata, 12; tenuifolia, 11, var. pseudoislandica, f. septentrionalis, 11, var. reticulata, 11; Tilesii, 12; *viridis*, 12
Chariomma nobilis, 729
Cheilanthes, 57; alabamensis, 57; *dealbata*, 61; Feei, 58; lanosa, 58; *lanuginosa*, 58; *vestita*, 58
Chisos Mountains, 794
Chlorine, relation of growth of orchid seedlings to, 371
Cladonia, 18; alpestris, 18; bacillaris, 18; cariosa, 18, f. cribrosa, 18; cenotea var. crossota, 18; chlorophaea, 19; cornuta, 18; crispata var. elegans, 18, var. infundibulifera, 18; cristatella, 18; deformis, 18; degenerans, 18; elongata, 18, var. ecmocyna, 18; fimbriata var. apolepta, 19, var. simplex f. major, 19, f. minor, 19; gracilis, 18, f. anthocephala, 18, var. dilatata, 18; pyxidata var. neglecta, 19; rangiferina, 18, f. nivea, 18; silvatica, 18, var. mitis, 18; subcervicornis f. turgescens, 19; unciialis, 18; verticillata, 18
Cladothrix lanuginosa, 796
Clematis alpina, 800
Coccidioides, 82; immitis, 85, 104
Coccocarpia, 441, 457; albida, 459, 466; asterella, 459; cronia, 458, 462, var. camporum, 458, var. erythrocardia, 458, var. granulosa, 458, 465, var. isidiophylla, 458, 463, var. lividorufa, 463, var. *primaria*, 463, var. prolificans, 458, 464; dominguensis, 459; elegans, 459, 466; epiphylla, 459; fibrillosa, 459; molybdaea, 457; *molybdaea* var. *cronia*, 461, 462, var. *tenuior*, 463; *parmelioides* var. *cronia*, 462; pellita, 458, 459, var. genuina, 458, var. lividorufa, 458, var. pannosa, 458, var. parmelioides, 458, 459, var. pyrrhichocarpa, 458, 460, var. strigosa, 458, 461; *pellita* var. *cronia*, 462, 463, var. *granulosa*, 465, var. *isidiophylla*, 463, var. *prolificans*, 464, var. *smaragdina*, 461, var. *tenuior*, 463; portoricensis, 458; subtilis, 458; tenuissima, 459
Cocoanut fiber, growth of orchid seedlings in, 577
Coenogoniaceae, 394
Coenogonium, 395; acrocephalum, 396; *andinum*, 396, 399; complexum, 397; confervoides, 396, 399; *controversum*, 398; depressum, 397, 402; dialeptizum, 397; disjunctum, 396; Echinus, 396; heterotrichum, 396, 400; implexum, 402; interplexum, 397, 401; interponendum, 397, 400; interpositum, 396, 401; Leprieurii, 396, 397, var. panniforme, 397; *Leprieurii* var. *subvirescens*, 397; Linkii, 396, 398; *Linkii* var. *Leprieurii* 397; moniliforme, 396, 397; pannosum, 397, 402; pulvinatum, 397; subvirescens, 396, 397; Tuckermani, 397
Collema, 404, 408; baculiferum, 409; subg. Blennothallia, 409; *bullatum*, 427, var. *dactylinoideum*, 427, var.

sertatum, 428; chilenum, 409; subg. Collemodiopsis, 408, 409; corynesporum, 409; crenatum, 409; cyrtaspis, 409, 412; *diaphanum*, 421; subg. **Dicollema**, 409, 411; glaucophthalmum, 409, 413, var. brasiliense, 409, var. granatense, 409, 413, var. typicum, 409; *glaucophthalmum*, 413; **Granadillae**, 408, 410; implicatum, 409, 414; Lactuca, 408; leptosporum, 409; leucocarpum, 409; leucopepla, 409, 411; lherminieri, 408, 410; mexicanum, 409; millegranum, 409; moluccanum, 436; nigrescens, 17, 409, var. caesium, 411; *nigrescens* var. *leucopepla*, 411; *oblique-peltatum*, 405; *olivaceum*, 424; *phyllocarpum*, 428; pycnocarpoides, 409; pycnocarpum, 409, 411, 412, var. **crassiusculum**, 409, 412, var. **Malmei**, 409, 412, var. **Minarum**, 409, 411, var. typicum, 409; *pycnocarpum*, 411, 412, f. *crassiusculum*, 412; **Ramboi**, 409, 415; saturninum, 422; solenarium, 409, stellatum, 409; subg. Synechoblastus, 409, 413; turgidulum, 409; *vesicatum*, 431

Collemaceae, 404

Coreopsis lanceolata, 797

Cornicularia divergens, 11

Costa Rica: botanical exploration in, 385; floristics of, 390; the foliose and fruticose lichens of, I, 373; geography of, 376; map of, *376*

Cotton plant, Super optimal and thermal death temperatures of, as affected by variations in relative humidity, 583

Crone's solution for growing orchids, 364

Cryptococcaceae, 104

Cryptococcus, 82, 474; *dermatitidis*, 85, 103; *gilchristi*, 85, 103; *hominis*, 85

Crystal violet, effect of, on growth of Endomyces capsulatus, 533

Cupressus arizonica, 796

Cyanisticta Hookeri var. **septentrionalis**, 17

Cystopteris, 58; bulbifera, 59, var. horizontalis, 60, *78;* fragilis, 58, 796

D

Davis Mountains, 793

Dennstaedtia, 60; punctilobula, 60

Dichodium, 407

Dicksonia, 60; *pilosiuscula*, 60; *punctilobula*, 60

Dicollema, 411

Diplazenia, 645; *Achrestogyne*, 677; *aciculiaris*, 682; *acuminata*, 703; *albaviridis*, 690; *alexicaca*, 729; *androsaemifolia*, 729; *angustifolia*, 710; *atropurpurea*, 724; *atroviolacea*, 724, *α. latifolia*, 724, *1. scandens*, 724, *2. suberecta*, 724, *β. cuneata*, 724, *γ. cordata*, 724; *bella*, 717; *boliviensis*, 716; *Buchtienii*, 730; *congesta*, 676; *crassinoda*, 703; *cuspidata*, 730; *eximia*, 706; *Fendleri*, 676; *fragrans*, 713, *α. oppositifolia*, 713, *β. ternatifolia*, 714; *Gardneriana*, 728, *β. grandiflora*, 728; *gentianoides*, 732, *α. velutina*, 732, *α.* longiloba*, 732, *β. glabra*, 733, *1. obovata*, 733, *2. ovata*, 733, *β.* longiloba*, 733, var. *Pohliana*, 733; *glabra*, 709; *illustris*, 728, *α. tomentosa*, 728, *a. rotundifolia*, 728, *1. hirsuta*, 728, *2. pubescens*, 728, *b. elliptica*, 728, *1. hirsuta*, 728, *2. pubescens*, 728, *c. oblongifolia*, 728, *f. pilosa*, 728, *β. glabra*, 729, f. *glabra*, 729; *linariaefolia*, 679; *linearis*, 733; *longiloba*, 736; *Luetzelburgii*, 698; *Martiana*, 702, *α. pubescens*, 702, *β. glabra*, 703; *mollis*, 730; *Moricandiana*, 705; *myriophyllum*, 682; *nobilis*, 729; *oblongifolia*, 708; *pastorum*, 679, var. *tenuifolia*, 680; *peduncularis*, 679; *pendula*, 725; *piladenia*, 730; *Pohliana*, 733; *polymorpha*, 679, *α. tenuifolia*, 679, *1. glabra*, 680, *2. volubilis*, 680, *3. puberula*, 680, *β. intermedia*, 680, *γ. peduncularis*, 680, *δ. brevifolia*, 680; *pulchella*, 736; *Riedelii*, 714; *Rosa-campestris*, 728; *rotundifolia*, 730; *sancta*, 726; *Sanderi*, 720; *saponaria*, 735; *Saponariae*, 734; *scabra*, 717; *Sellowii*, 719; *spigeliaeflora*, 736, *β. longiloba*, 736; *splendens*, 707; *stenoloba*, 676; *surinamensis*, 704; *tenuifolia*, 679, 680, *β. puberula*, 679, *γ. volubilis*, 679, f. *pastorum*, 680; *tetradenia*, 730; *upatae*, 704; *urophylla*, 722; *velutina*, 732; *venulosa*, 723; *vincaeflora*, 679; *xanthostoma*, 734, *a. major*, 734, *b. minor*, 734, f. *longiloba*, 736

Diploschistes bryophiloides, 20; **canadensis**, 20

Dodge, Carroll W. The foliose and fruticose lichens of Costa Rica. I, 373

Drymaria gracilis, 797

Dryopteris, 66; *acrostichoides*, 63

Dyes: effect of various, on growth of Endomyces capsulatus, 523

E

Echites, 629, 646, 777; *acuminata*, 635, 744; *acutiloba*, 649; *alexicaca*, 729; *almadensis*, 763; *altescandens*, 741; *andina*, 665; *Andrieuxii*, 688; *angustifolia*, 640, 774; *antennacea*, 741; *apocynifolia*, 661; *arenaria*, 697; *assimilis*, 665; *atropurpurea*, 724; *atroviolacea*, 724; *augusta*, 789; *auriculata*, 763; *Benthamii*, 774; *bicorniculata*, 634; *bogotensis*, 677; *Boliviana*, 741; *brachyloba* 670; *Brachysiphon*, 780;

brachystachya, 752; *bracteata*, 673, 761; *bracteosa*, 742; *breviflora*, 640; *brevipes*, 638; *Brownei*, 656; *campestris*, 762; *ciliata*, 763; *citrifolia*, 637; *Cobanensis*, 648; *coccinea*, 734, β. *ovata*, 735; *comosa*, 747; *congesta*, 675; *convolvulacea*, 687; *cordata*, 686; *Coulteri*, 657; *crassinoda*, 703; *cuspidata*, 630; *cuspidifera*, 744; *Cuyabensis*, 752; *densiflora*, 668; *dichotoma*, 644; *disadena*, 630; *emarginata*, 672; *erecta*, 672; *Fluminensis*, 763, β. *Claussenii*, 763; *fragrans*, 713; *Franciscea* var. *pallidiflora*, 764; *funiformis*, 697; *Ganabarica*, 697; *glandulosa*, 692; *glaucescens*, 686; *gracilis*, 743; *grandiflora*, 789, var. *major*, 789, var. *minor*, 787; *Guarantica*, 789; *Guayaquilensis*, 743; *Guianensis*, 740; *hirsuta*, 749, 762, α. *latifolia*, 763, β. *angustifolia*, 750; *hirtella*, 686, 743; *hirtiflora*, 744; *hispida*, 763; *hypoleuca*, 781; *illustris*, 727; *japurensis*, 630; *jasminiflora*, 743; *javitensis*, 775; *Karwinskii*, 657; *Lambertiana*, 787; *lanuginosa*, 782; *lasiocarpa*, 750, β. *angustifolia*, 750, γ. *Lobbiana*, 750; *laxa*, 695; *leptophylla*, 756; *linearifolia*, 640, 756; *longiflora*, 789; *macrophylla*, 678, 762; *Macrosiphon*, 783; *Mansoana*, 636; *Maranhamensis*, 752; *Martiana*, 702; *membranacea*, 744; *mexicana*, 660; *microcalyx*, 744, β. *glabra*, 744; *microphylla*, 697; *minima*, 643; *mollissima*, 751; *montana*, 664; *mucronata*, 743; *multiflora*, 788; *myrtifolia*, 642; *Oaxacana*, 686; *obovata*, 705; *ovalis*, 626; *pallida*, 631; *palustris*, 763; *pastorum*, 679; *Pavonii*, 749; *pedunculariis*, 679; *petraea*, 787; *pinifolia*, 787; *Pohliana*, 732, var. α. *angustifolia*, 732, var. β. *latifolia*, 732; *Prieurei*, 740; *pubescens*, 676; *pubiflora*, 752; *pulchella*, 736; *pycnantha*, 668; *pycnanthe*, 668; *repens*, 639; *Richardi*, 762; *rigida*, 631; *riparia*, 665; *Rosa-campestris*, 728; *Rosana*, 652; *rosea*, 642; *rugosa*, 754; *sancta*, 726; *Sanctae-Crucis*, 634; *scabra*, 752; *secundiflora*, 744; *Smithii*, 660; *speciosa*, 743; *spigeliaeflora*, 736; *splendens*, 707; *Stadelmeyeri*, 763; *suaveolens*, 695, 781; *subcarnosa*, 772; *subsagittata*, 743; *subsessilis*, 667; *subspicata*, 739; *surinamensis*, 630; *symphitocarpa*, 755; *tenuicaulis*, 752; *tenuifolia*, 679; *tomentosa*, 762, β. *laticordata*, 763; *torosa*, 656, var. *Brownei*, 656; *torulosa*, 642, 656; *trifida*, 630, 631, f. *puberula*, 635, var. *Sanctae Crucis*, 634; *tubiflora*, 648; *tubulosa*, 630; *Velame*, 788; *velutina*, 732; *venenosa*, 728; *Veraguasensis*, 690; *versicolor*, 752; *virescens*, 784, 785; *xanthostoma*, 734

Echitoideae, The American genera of, of Apocynaceae, 605
Eclipta integrifolia, 34
Endomyces capsulatus Rewbridge, Dodge & Ayers: a causative agent of fatal cerebrospinal meningitis, 471; animal inoculations, 543; carbohydrate reactions, 510; cultural characteristics, 494; cytology, 477; dyes, reaction to, 523; gelatine liquefaction, 519; historical review, 472; hydrogen sulphide production, 519; indol and skatol reactions, 517; light reactions, 520; nitrates, reduction of, 519; oxygen, free, relation to, 518
Endomyces, 82, 103; capsulatus, 471, *554*, *556*, *558*, *560*, *562*, *566*, *568*, var. isabellinus, 472, 473, *564*; dermatitidis, 106, *116*, 472
Endomycetaceae, 104
Eosine B, effect of, on growth of Endomyces capsulatus, 535
Ephebaceae, 402
Ephebia, 403
Equisetum, 69; arvense, 69; *hyemale* var. *robustum*, 70; laevigatum, 71; praealtum, 70; *robustum*, 70
Eriadenia, 645; *obovata*, 771
Erioderma, 440, 455; Leylandi, 455; limbatum, 455, 456; microcarpa, 455; *physcioides*, 451; polycarpum, 455; pulchrum, 455; verruculosum, 455; Wrightii, 455; *Wrightii* var. *limbatum*, 456
Eriogonum Havardii, 800; pannosum, 800
Eriogynia caespitosa, 800
Eupatorium Parryi, 797
Euphorbia eriantha, 798
Exadenus, 138; *alatus*, 139; *brevicornis*, 140; *parviflorus*, 140, var. β. *latifolius*, 142; *paucifolius*, 140
Exothostemon, 645; *bracteatum*, 761; *gracile*, 743; *Javitense*, 775; *macrophyllum*, 762; *mollissimum*, 751; *speciosum*, 743

F

Fernaldia pandurata, 777
Ferns and fern allies of Missouri, 45
Festuca ligulata, 801
Filix, 59; *bulbifera*, 59; *fragilis*, 59
Frasera speciosa, 798
Fructose, growth of orchid seedlings in solutions containing, 574, 575

G

Gabura, 408
Galactose, growth of orchid seedlings in solutions containing, 574, 575

Gelatine liquefaction in cultures of Endomyces capsulatus, 519
Gentiana viridis, 151
Geranium caespitosum, 798
Germination and growth of orchid seedlings, sources of carbohydrate for, 569
Glucose, growth of orchid seedlings in solutions containing, 574, 579
Gonionema, 403; *velutina*, 403
Graphis scripta, 20
Guadalupe Mountains, 794

H

Haemadictyon glandulosum, 692
Halenia, A monograph of the American species of the genus Halenia, 119
Halenia, 137; **adpressa**, 148, *218;* alata, 139, *214;* *antigonorrhoica*, 202; *apiculata*, 178; asclepiadea, 205, *216;* *asclepiadea*, 188; barbicaulis, 198, *220;* bella, 206, *220;* **bifida**, 201, *216;* *Brentoniana*, 167; brevicornis, 140, 147, *214*, var. **chihuahuensis**, 144, *214*, var. **divergens**, 144, *214*, var. **latifolia**, 141, *214*, var. **micranthella**, 143, var. **multiflora**, 142, *214*, var. **ovata**, 145, *214*, var. **Tuerckheimii**, 145, *214;* caespitosa, 184, *216;* **caleoides**, 173, *214;* *candida*, 161; *chlorantha*, 172; Conzattii, 171, *214;* crassiuscula, 159, *214;* dasyantha, 154, *218;* decumbens, 178, *214;* deflexa, 162, *214*, *222*, var. Brentoniana, 167; *Dombeyana*, 197, var. *α.* 194, 196, var. *brevicornis*, 182; elata, 155, *220;* **elegans**, 203, *218;* *elongata*, 175; *erythraeoides*, 140; foliosa, 153, *220;* gentianoides, 152, *218;* **gigantea**, 192, *218;* gracilis, 186, *216*, var. *Dombeyana*, 197; guatemalensis, 179, *214*, var. **latifolia**, 180; sect. Haleniastrum, 158; Herzogii, 195; *heterantha*, 162; Hieronymi, 201; Hoppii, 204, *216;* hygrophila, 156, *218;* hypericoides, 184; inaequalis, 150, *218;* *inaequalis*, 151; *Jamesoni*, 186; Kalbreyeri, 206, *218;* Karstenii, 149, *218;* **Killipii**, 187, *216;* *longicornu*, 178; macrantha, 148; major, 158, *218;* Mathewsii, 188; Meyeri Johannis, 197, *216;* *Michauxiana*, 162; *micranthella*, 143; **minima**, 193, *216;* *multiflora*, 142; nudicaulis, 174, *214;* *nutans*, 175; Palmeri, 171, *214;* **parallela**, 157, *218;* *parviflora*, 140, var. *latifolia*, 142; *paucifolia*, 140; *Pavoniana*, 197; penduliflora, 194, *216;* phyteumoides, 194; *pichinchensis*, 186; pinifolia, 185; plantaginea, 175, *214*, f. **grandiflora**, 176, var. *latifolia*, 180; **platyphylla**, 173, *214;* Pringlei, 160, *214;* pulchella, 185, *220;* Purdieana, 199, *218*, var. **congesta**, 200, *216;* *Purpusi*, 174; pusilla, 182, *216;* **recurva**, 161, *214;* *reflexa*, 162; **rhyacophila**, 168, *214*, var. **macropoda**, 170, var. **procumbens**, 170; robusta, 190, *216;* *Rothrockii*, 161; Rusbyi, 199, *216;* *scapiformis*, 174; Schiedeana, 172, *214;* **Schultzei**, 151; Shannonii, 177, *214*, f. **compacta**, 178; silenoides, 195, *216;* **spatulata**, 183, *216;* sphagnicola, 207; stellarioides, 152; Stuebelii, 190, *216;* subinvolucrata, 149, *220;* sect. Swertiella, 139; taruga gasso, 191, *218;* Tolimae, 155, *218;* *Tuerkheimii*, 145; umbellata, 196, *216;* valerianoides, 182; verticillata, 153, *218;* vincetoxicoides, 188, *216;* viridis, 151, *220;* **Weberbaueri**, 189, *216;* Weddelliana, 202, *216*
Helleranthus quadrangulatus, 313
Heterothrix, 645; *pycnantha*, 668; *Vanheurckii*, 771
Hieracium carneum, 798
Hippochaete prealta, 70
Holocoenis Leprieurii, 397
Holodiscus dumosus, 800
Homaladenia, 645; *brevifolia*, 680; *linariaefolia*, 680; *pastorum*, 680; *pedunculatis*, 680; *puberula*, 680; *tenuifolia*, 680; *vincaeflora*, 680
Humidity, relative, Super optimal and thermal death temperatures of the cotton plant as affected by variations in, 583
Hydrogen ion concentrations: of nutrient solutions in relation to growth of orchids, 227, 579; various, reaction of Endomyces capsulatus to, 502
Hydrogen sulphide production in Endomyces capsulatus, 519
Hymenopappus radiatus, 798
Hypopitys sanguinea, 799

I

Indol, reaction of, to Endomyces capsulatus, 517
Iron in nutrient solutions for orchids, 226, 365, 572
Isoetes, 74; Butleri, 75; Engelmanni, 74; melanopoda, 75

J

Juniperus flaccida, 796

K

Keerlia, 37; bellidifolia, 37; effusa, 38; *linearifolia*, 33; *mexicana*, 31; *ramosa*, 24, 25; *skirrhobasis*, 24, 25
Knudson's solution for growing orchids, 365, 572, 574
Kujala, Dr. Viljo, lichens collected by, 7

L

La Garde's solution for growing orchids, 226, 234, 363, 572, 578
Laphamia quinqueflora, 798
Lappula grisea, 797
Larsen, Esther L. Astranthium and related genera, 23
Laseguea, 645; *acutifolia*, 673, f. *α. Guilleminiana*, 674, *1. Griseo-olivacea*, 674, *2. Griseo-fusca*, 674, f. *β. obliquinervia*, 674, *αα. ovata*, 674, *1. griseo-olivacea*, 674, *ββ. obovata*, 674, f. *γ. Glabrescens*, 674, f. *δ. scabrinervis*, 674, f. *ε. glabra*, 674; *antennacea*, 741; *bracteata*, 671; *emarginata*, 673; *erecta*, 673, *α. Guilleminiana*, 673, *1. griseo-olivacea*, 673, *2. griseo-fusca*, 673, *β. obliquinervia*, 673, *αα. ovata*, 673, *1. griseo-olivacea*, 673, *ββ. obovata*, 673, *γ. glabrescens*, 673, *δ. acutifolia*, 673, *ε. scabrinervia*, 673, *ζ. glabra*, 673; *foliosa*, 658; *glabra*, 673; *Guilleminiana*, 672; *Hookeri*, 671; *Mandoni*, 671; *obliquinervia*, 673; *Pentlandiana*, 671; *subspicata*, 740; *venustula*, 740; *villosa*, 747
Lastrea, 66
Lathargium, 413
Lecania dimera, 14
Lecanora coilocarpa, 14; subintricata, 14; symmicta, 14; umbrina, 14
Lecidea goniophila, 19; lapicida, 19; *parmelioides*, 459
Lemmopsis polyschidioides, 404
Lempholemma, 404; compactum, 404; **dichotomum**, 406; Dussii, 405; **obliquepeltatum**, 405
Lepidocollema carassense, 440
Leprocollema americanum, 404; Finkii, 404
Leptogidium byssoides, 403
Leptogium, 404, 416; adpressum, 418; austroamericanum, 419; azureum, 420, 433; brasiliense, 420; *bullatum*, 426; callithamnium, 418; chloromelan, 419; *chloromelan*, 420; conchatum, 420; coralloideum, 419, 430; *corrugatulum*, 431; denticulatum, 419, 432; diaphanum, 418, 421; *diaphanum* f. *coralloideum*, 430; dimorphum, 420; subg. Diplothallus, 418, 420; subg. Euleptogium, 418, 424; foveolatum, 418, 425; fusisporum, 418; granulare, 419; *Hildenbrandii* var. *papillosum*, 422; *hypotrachynum*, 419; inflexum, 418, 423, var. isidiosulum, 418, 423; laevius, 420; Lafayetteanum, 420; *leptophyllinum*, 432; subg. Mallotium, 418, 422; marginellum, 408, 419, 431; mattogrossense, 419; megapotamicum, 418; microcarpum, 419; microstictum, 418, 426; *moluccanum*, 436; olivaceum, 418, 424, var. **granulosum**, 424; pachycheilum, 420; **papillosum**, 418, 422; phyllocarpum, 419, 428, var. campestre, 419, 430, var. macrocarpum, 419, 429; *phyllocarpum* var. *coralloideum*, 430, var. *isidiosum*, 430; pichneum, 419; pilcomayense, 419; Puiggarii, 420; pulchellum, 420; *punctulatum*, 421; resupinans, 418; reticulatum, 418; saturninum, 17; Schiffneri, 420; sessile, 419; simplicius, 420, 437, var. **pichneoides**, 419, 438; **Standleyi**, 420, 435; stellans, 420; stipitatum, 418, 428; tremelloides, 17, 416; *tremelloides*, 435, var. *azureum* f. *isidiosum*, 432, var. *impressopunctatum*, 421, var. *leptophyllinum*, 432, var. *minor*, 436; **Tuckermani**, 420, 436; ulvaceum, 420; vesiculosum, 418, 426, var. digitatum, 427
Letharia vulpina, 11, f. compacta, 11
Lichen flora of North America, Contribution to the, 7
Lichen azureum, 434; *bullatum*, 426; *crispus*, 408; *diaphanum*, 421; *marginellus*, 431; *melanocarpus*, 394; *pannosus*, 453; *velutinus*, 403; *vesiculosus*, 426
Lichens of Costa Rica, The foliose and fruticose, 373
Light: white, effect of, on growth of Endomyces capsulatus, 521; of red, 522
Linum Schiedeanum, 799
Lithospermum viride, 797
Lobaria linita, 17; oregana, 17; pulmonaria, 17
Lobarina verrucosa, 17
Loeselia Greggii, 799
Lonicera arizonica, 797
Lorinseria areolata, 69
Lycopodium, 72; complanatum flabelliforme, 73; lucidulum, 73, var. porophilum, 73; *porophilum*, 73

M

Macrosiphonia, 777; *Balansae*, 788; *Berlandieri*, 783; Brachysiphon, 780; subgen. **Eumacrosiphonia**, 784; *Guarantica*, 789; Hesperia, 779; hypoleuca, 781; lanuginosa, 782; longiflora, 789; *longiflora* var. *Guarantica*, 790; Macrosiphon, 783; Martii, 784; *pedunculoris*, 788; petraea, 786, var. **pinifolia**, 787, 788, var. typica, 787; *pinifolia*, 788, f. *glabrata*, 788, var. *intermedia*, 788, f. *pedunculoris*, 788, f. *setosa*, 788; *prostrata*, 787; subg. **Telosiphonia**, 778; Velame, 788; *Velame β. goyazensis*, 788; *verticillata*, 786, *α. petraea*, 787, *β. intermedia*, 787, *γ. pedunculoris*, 787, *δ. pinifolia*, 787; *virescens*, 785, var. *Missionum*, 785; *Wrightii*, 781

Magnesium, relation of growth of orchid seedlings to, 231, 366
Mahonia repens, 796
Mallotium, 422
Malmella, 440, 447; **albida**, 448, 449; **caesiocinerea**, 448, 452; **physcioides**, 448, 451; **Randii**, 448; **Santamariae**, 448, 450
Maltose, growth of orchid seedlings in solutions containing, 226, 574, 575
Mandevilla, 645; **Achrestogyne**, 677; acutiloba, 649; albo-viridis, 690; anceps, 774; Andrieuxii, 645, 688; **angustifolia**, 710; antennacea, 741; apocynifolia, 660; **atroviolacea**, 724; *attenuata*, 761; *auriculata*, 764; *Bangii*, 693; **bella**, 717; Benthamii, 644, 774; **bogotensis**, 677; *Boliviana*, 741; **boliviensis**, 716; brachyloba, 670; bracteata, 761; **bracteosa**, 742; Bridgesii, 693; callacatensis, 663; **callista**, 700; **caurensis**, 770; cercophylla, 669; **cereola**, 712; **coccinea**, 734; **congesta**, 675; convolvulacea, 687; **crassinoda**, 703; **cuspidata**, 730; *denticulata*, 764; *Dielsiana*, 772; Donnell-Smithii, 650; equatorialis, 689; erecta, 672; subg. **Eumandevilla**, 647; **eximia**, 706; subg. **Exothostemon**, 737; Fendleri, 758; *fluminensis*, 764; foliosa, 658; fragilis, 667; **fragrans**, 713; funiformis, 697; *funiformis* var. *peduncularis*, 697; **glabra**, 709; *glabra*, 771; glandulosa, 692; grata, 696; hirsuta, 762; *hispida*, 764; **illustris**, 727, var. **glabra**, 729, var. typica, 727; **immaculata**, 721; Jamesonii, 666; javitensis, 775; Karwinskii, 657; lancifolia, 773; lasiocarpa, 750; laxa, 695; sect. laxae, 682; leptophylla, 756; linearis, 733; *linearis*, 756; *Loesneriana*, 690; **lucida**, 718; **Luetzelburgii**, 698; *Mandoni*, 693; **Martiana**, 702, var. **glabra**, 703, var. typica, 702; mexicana, 660; **minor**, 711; mollissima, 751; montana, 664; *montana* var. *peruviana*, 665; sect. montanae, 661; **Moricandiana**, 705, var. **bahiensis**, 706, var. typica, 705; Moritziana, 768; **Muelleri**, 717; **myriophyllum**, 682; oaxacana, 644, 686; **oblongifolia**, 708; *palustris*, 764; *parvifolia*, 753; Pavonii, 749; **pendula**, 725; Pentlandiana, 671; **permixta**, 715; platydactyla, 651; polyantha, 769; *polyantha*, 769; *potosina*, 777; pycnantha, 668; riparia, 665; Rosana, 652; rugosa, 754; *Rusbyi*, 764; rutila, 757; sagittarii, 768; **sancta**, 726; **Sanderi**, 720; scaberula, 761; scabra, 752; Schlimi, 759; *Schlimii*, 759; *Schumanniana*, 686; **scutifolia**, 663; **Sellowii**, 719; sertuligera, 649, 654; **spigeliaeflora**, 736; **splendens**, 707; Spruceana, 776; *suaveolens*, 695; **subcarnosa**, 644, 772; *subcarnosa*, 772; subcordata, 693; subpaniculata, 678; subsagittata, 644, 645, 743; subscorpioidea, 653; subsessilis, 667; subspicata, 645, 739; superba, 709; **surinamensis**, 704; symphitocarpa, 755; Syrinx, 649, 653; *tenuicarpa*, 741; tenuifolia, 679; sect. tenuifoliae, 679; *tomentosa*, 764, var. *hirsuta*, 764, var. *hispida*, 764, var. *Vahleana*, 764; torosa, 644, 656; sect. torosae, 655; Trianae, 760; tubiflora, 648; sect. tubiflorae, 647; *Tweediana*, 695; Ulei, 771; **urophylla**, 722; Vanheurckii, 771; **velutina**, 731, var. glabra, 732, var. typica, 732; *velutina*, 777; **venulosa**, 723; veraguasensis, 690; *veraguensis*, 690; villosa, 747
Manganese sulphate in nutrient solutions, 226
Mannose, growth of orchid seedlings in solutions containing, 574, 577
Matteuccia nodulosa, 65; *Struthiopteris*, 65
Media, growth of Endomyces capsulatus in various, 494
Meningitis, fatal cerebrospinal, Endomyces capsulatus as a causative agent of, 471
Mentzelia asperula, 799
Mesechites, 629; acuminata, 635; *Andrieuxii*, 645, 688; *angustata*, 644, 775; angustifolia, 640; bicorniculata, 634; *brevipes*, 638; *Brownei*, 644, 656; citrifolia, 637; dichotoma, 644; subg. **Didymadenia**, 638; *disadena*, 631; subg. **Eumesechites**, 630; *Guayaquilensis*, 644, 744; *Guianensis*, 645, 740; *hastata*, 644, 744; *hirtella*, 644, 744; *hirtellula*, 644, 686; *japurensis*, 630; *jasminiflora*, 644, 744; *lanceolata*, 639; *linearifolia*, 640; **Mansoana**, 636; minima, 643; *myrtifolia*, 642; *Oaxacana*, 644, 686; repens, 639; rosea, 641; Sanctae-Crucis, 634; *subcarnosa*, 644, 772; *sulphurea*, 636; *surinamensis*, 631; *torulosa*, 644, 656; trifida, 630, 644
Methylene blue, effect of, on growth of Endomyces capsulatus, 534
Micradenia, 645; *atroviolacea*, 724; *crassinoda*, 703; *fragrans*, 714; *hirsutula*, 702; *Martiana*, 702; *Moricandiana*, 706; *Riedelii*, 740; *Sellowii*, 719; *splendens*, 707; *urophylla*, 722
Missouri, Ferns and fern allies of, 45; A neutral (?) strain of Mucor sphaerosporus from, 469
Mitozus brachystachyus, 753; *brevipes*, 638; *Cuyabensis*, 753; *exilis*, 697; *funiformis*, 697; *Guanabaricus*, 697; *leptophyllus*, 756; *microphylla*, 697; *rugosus*, 755; *symphitocarpus*, 755; *tenuicaulis*, 753; *versicolor*, 752

Monilia, 82, 103
Moore, John Adam, Steyermark, Julian A. and. Report of a botanical expedition into the mountains of western Texas, 791
Moore, Morris. Blastomycosis: report of a case, with a study of an etiologic factor and a classification of the organism, 79; A neutral (?) strain of Mucor sphaerosporus from Missouri, 469; A study of Endomyces capsulatus Rewbridge, Dodge and Ayers: a causative agent of fatal cerebrospinal meningitis, 471
Mucor sphaerosporus from Missouri, A neutral (?) strain of, 469
Mycoblastus melinus, 19; sanguinarius, 19, var. **Dodgeanus**, 19
Mycoderma, 474
Mycoderma dermatitis, 103

N

Nama Havardii, 799; xylopodum, 799
Nephrodium, 66
Nephromium **canadense**, 17; laevigatum var. papyraceum, 17; subparile, 16
Nephromopsis ciliaris, 13; platyphylla, 13
Nerium sarmentosum, 642
Nitrates, reduction of, in Endomyces capsulatus, 519
Nitrogen in relation to growth of orchid seedlings, 234, 368.
Nothochlaena pulchella, 61
Notholaena, 60; dealbata, 60; Greggii, 796; *nivea dealbata*, 61
Nutrient solutions for orchids, 225, 363, 571
Nutritional ions, The sensitivity of orchid seedlings to, 223
Nymphaea, A new yellow, from tropical Africa, 2
Nymphaea (§ Brachyceras) **Burttii**, 1, *6*; Stuhlmannii, 3; sulphurea, 3

O

Obletia, 251
Ochrolechia upsaliensis, 14
Odontadenia cuspidata, 730; *glandulosa*, 692
Oidium, 82, 102, 474
Onoclea, 61; sensibilis, 61; *Struthiopteris*, 64
Oosporaceae, 104
Ophioglossum, 46; Engelmanni, 47; vulgatum, 46
Orbilia, 396
Orchid seedlings, Nutrient solutions for, 363; The sensitivity of, to nutritional ions, 223; Sources of carbohydrate for germination and growth of, 569
Osmotic concentration, relation of, to growth of orchid seedlings, 231
Osmunda, 71; cinnamomea, 72; Claytoniana, 72; regalis spectabilis, 71; *Struthiopteris*, 64
Ostrya Baileyi, 797
Oxygen, free, relation of Endomyces capsulatus to, 518

P

Pannaria, 440, 441; applanata, 441; brasiliensis, 442; caeruleo-nigricans, 442; imbricatula, 442; isidioidea, 442, var. **pulvinata**, 443; Malmei, 441; *Mariana* var. *isidioidea*, 442, f. *radiata*, 446; **Moseni**, 442, 443; **radiata**, 442, 446; rubiginosa, 441; *rubiginosa*, 443, 445, f. *caesiocinerea*, 452, f. *cinerascens*, 452; stylophora, 442, 445; **Vainii**, 442, 445
Pannariaceae, 438
Parmelia ambigua, 13; *cronia*, 462; Delavayi, 13; *diaphana*, 421; elegantula var. **americana**, 14; enteromorpha, 13; *exasperatula*, 13; *hyperopta*, 13; *lividorufa*, 463; lophyrea, 13; *molliuscula*, 13; pallescens, 13; papulosa, 13; *pellita*, 459; physodes f. labrosa, 13, f. vittata, 13; pubescens var. reticulata, 14; subaurifera, 13; sulcata, 13; vagans, 13; vittata, 13
Parmeliella, 440, 452; *corallinoides*, 453; microphylloides, 453; miradorensis, 453; nigrocincta, 453, var. Weddellii, 453; pannosa, 453, var. **coralloidea**, 454; triptophylla, 453
Parsonsia bracteata, 671
Patinella, 396
Pellaea, 61; atropurpurea, 62, var. *Bushii*, 62, var. cristata, 62; *dealbata*, 61; glabella, 62
Peltigera aphthosa, 16; canina, 16, var. membranacea, 16, var. suomensis, 16; horizontalis, 16; lepidophora, 16; malacea, 16; Maurizii, 16; microphylla, 16; Nylanderi, 16; *perfida*, 16; rufescens, 16; *suomemsis*, 16; variolosa, 16; venosa, 16
Pentoses, growth of orchid seedlings in, 576
Pentstemon baccharifolius, 801
Perityle Parryi, 798
Perry, Lily M. A revision of the North American species of Verbena, 239
Pertusaria *amara*, 14; faginea, 14; multipuncta, 14; pertusa, 14; xanthostoma, 14
Phegopteris, 66; *hexagonoptera*, 68
Phlyctis argena, 14
Phosphorus, growth of orchids in solution containing, 233, 367
Phryma caroliniensis, 309
Physcia, 15; *adglutinata* var. *pyrithro-*

cardia, 15; *astroidea*, 15; caesia, 15; Clementiana, 15; endophoenicea, 15; grisea var. semifarrea, 15; muscigena, 15, f. lenta, 15, f. squarrosa, 15; pyrithrocardia, 15; stellaris, 15, var. rosulata, 15
Physma, 404; Boryanum, 407; byrsinum, 408; pruinosum, 408
Pilophoron aciculare, 17
Pinaropappus parvus, 798
Pinkerton, M. Elizabeth. Ferns and fern allies of Missouri, 45
Placodium chrysodetum, 15; citrinum, 15; elegans, 15; ferrugineum, 15; xanthostigmoideum, 15
Platyema septentrionale, 12
Polychidium, 403
Polygala rimulicola, 800
Polypodium, 62; *incanum*, 63; polypodioides, 63; virginianum, 63; *vulgare*, 63
Polystichum, 63, 66; acrostichoides, 63, var. incisum, 64, var. *Schweinitzii*, 64
Polystichum, 63
Potamogeton clystocarpus, 802; diversifolius, 802
Potassium in relation to growth of orchid seedlings, 231, 234, 361
Prestonia hirsuta, 749; *peruviana*, 692
Prestoniopsis Fendleri, 676; *hirsuta*, 676; *pubescens*, 676; *venosa*, 676
Pring, George H. and Robert E. Woodson, Jr. A new yellow Nymphaea from tropical Africa, 1
Prunus Havardii, 800
Psora decipiens, 19
Psoroma cinchonarum, 440
Pteretis, 64; nodulosa, 64
Pteridium, 65; *aquilinum*, 65; latiusculum, 65, var. pseudocaudatum, 65
Pteridophyta, new or rare, in Texas, 796
Pterigyopsis atra, 403
Pteris aquilina, 65
Pterospora Andromeda, 799

Q

Quercus hypoleuca, 798

R

Raffinose, growth of orchid seedlings in solutions containing, 574
Ramalina calicaris, 11; geniculata, 11; Roesleri, 11; thrausta, 11
Räsänen, Veli J. P. B. Contribution to the lichen flora of North America, 7
Report of a botanical expedition into the mountains of western Texas, 791
Rhabdadenia campestris, 763
Rhamnose, growth of orchid seedlings in solutions containing, 574, 576
Rhamnus fasciculata, 800
Rhodocalyx coccineus, 734; *hypoleucus*, 781; *lanuginosus*, 782; *ovatus*, 735; *suaveolens*, 781; *Tweedianus*, 735
Ribes mescalerium, 800
Riella americana, 795
Rinodina demissa, 15
Rio Grande, canyons of the, 794
Robinia luxurians, 799
Roccellaceae, 394
Rosa mirifica, 800

S

Saccharomycetaceae, 104
Saccharomyces, 82, 103, 474; *hominis*, 103
Salep, 577
Salt per cent, relation of growth of orchid seedlings to, 371
Scytenium, 408
Secondatia stans, 659
Selaginella, 74; apoda, 74; *apus*, 74; Pringlei, 796; rupestris, 74
Sensitivity of orchid seedlings to nutritional ions, 223
Shive's solution for growing orchids, 572, 575
Shuttleworthia, 251
Siphula ceratites, 14
Sisymbrium diffusum, 798
Skatol, reaction of, to Endomyces capsulatus, 517
Sodium borate in nutrient solutions, 226
Solidago Wrightii, 798
Solorina saccata, 17
Sorghastrum nutans, 801
Spermatophyta, new or rare, in Texas, 796
Sphaerophoraceae, 393
Sphaerophorus globosus var. lacunosus, 20; melanocarpus, 394; Tuckermanii, 20
Sphenopholis obtusata, 801
Spilanthes, 23
Spiranthes saltensis, 802
Sporotrichum, 82
Squamaria alphoplaca, 14
Staurothele circinata, 20
Stephanophorus, 424
Stereocaulon tomentosum, 14
Steyermark, Julian A., John Adam Moore and. Report of a botanical expedition into the mountains of western Texas, 791
Sticta scrobiculata, 17
Stictina fuliginosa, 17
Struthiopteris germanica, 65
Study Butte area, 794
Styleurodon carolinianum, 309
Sucrose, growth of orchid seedlings in solutions containing, 574, 576
Sugars, growth of orchid seedlings in solutions containing various, 226, 571
Sulphate, relation of growth of orchid seedlings to, 368

Super optimal and thermal death temperatures of the cotton plant as affected by variations in relative humidity, 583
Swertia, 138; *americana*, 162; *asclepiadea*, 205; *brevicornis*, 140; *corniculata*, 162; *cucullata*, 140; *deflexa*, 162; *gracilis*, 186; *hypericoides*, 184; *Michauxiana*, 162, 172; *parviflora*, 140, var. *angustifolia*, 140, var. *latifolia*, 142; *pinifolia*, 185; *plantaginea*, 175; *quadricornis*, 205; *recurva*, 161; *umbellata*, 191
Symphoricarpos rotundifolius, 797
Synechoblastus, 413

T

Temnadenia, 646; *cordata*, 686, *glaucescens*, 686; *lasiocarpa*, 750; *Lobbiana*, 750; *pallidiflora*, 764; *palustris*, 764; *secundiflora*, 744; *tomentosa*, 764; *xanthostoma*, 735
Temperature chamber for treating cotton plants, 585, *586*
Temperatures: reaction of Endomyces capsulatus to, 500; Super optimal and thermal death, of the cotton plant as affected by variations in relative humidity, 583
Tetragonanthus, 138; *alatus*, 139; *Brentonianus*, 167; *brevicornis*, 140; *decumbens*, 178; *deflexus*, 163, var. *Brentonianus*, 167, var. *heteranthus*, 163; *heterantherus*, 163; *heteranthus*, 163; *longicornis*, 178; *Palmeri*, 171; *parviflorus*, 140; *paucifolius*, 140; *plantagineus*, 175; *Rothrockii*, 161; *Schiedeanus*, 172; *Whitei*, 188
Texas: Report of a botanical expedition into the mountains of western, 791; species new or rare in the flora of, 795, additional records of interest, 802
Thamnolia vermicularis, 14
Thelypteris, 66; hexagonoptera, 68; marginalis, 67; palustris pubescens, 66; spinulosa, 67, var. intermedia, 67; *Thelypteris*, 66
Thermutis, 403; velutina, 403
Tillandsia recurvata, 801
Trachelopsermum stans, 659
Tradescantia Wrightii, 801
Trentepohlia, 396

U

Usnea caucasica, 9; comosa, 9, var. stuppea, 9; dasypoga, 9, var. subscabrata, 9; hirta, 9; Kujalae, 9; lapponica, 9; longissima, 9; prostrata, 9; rugulosa, 9; similis, 9; *subcomosa*, 9
Uwarowia, 251

V

Valeriana texana, 801
Vauquelinia angustifolia, 800
Venturiella sinensis, 796
Verbena, A revision of the North American species of, 239
Verbena, 251; *affinis*, 257; *Alopecurus*, 285; *ambrosifolia*, 326, f. eglandulosa, 328; amoena, 341; Andrieuxii, 333; *angustifolia*, 282; *arizonica*, 300, 337; *Aubletia*, 315, var. *Drummondi*, 316; Aubletia var. Lamberti, 343; barbata, 342; bipinnatifida, 323, var. latilobata, 325; *biserrata*, 268; bonariensis, 254; *bonariensis* var. *littoralis*, 257, var. *rigida*, 253; *bracteata*, 304, *358;* bracteosa, 304, var. *albiflora*, 304, var. *brevibracteata*, 304; brasiliensis, 255; *brevibracteata*, 338; canadensis, 315, *358;* canadensis var. *Ehrenbergii*, 318, subsp. *elegans*, 318, var. *Lamberti*, 316; canescens, 301, var. Roemeriana, 302; *canescens*, 304, var. *neomexicana*, 296; *caracasana*, 257; carnea, 309; carolina, 268; *caroliniana*, 268, 309, f. or var. *polystachya*, 268, f. or var. *recta*, 271; *carolinensis*, 309; ciliata, 330, var. longidentata, 331, var. pubera, 332; *confinis*, 304; *cuneifolia*, 286; delicatula, 342; delticola, 314; *diffusa*, 275; *domingensis*, 262; *Drummondii*, 316; Ehrenbergiana, 267; elegans, 318, var. asperata, 319; *elongata*, 254; erinoides, 341; *exilis*, 334; Gooddingii, 335, var. nepetifolia, 337; gracilis, 300; *grandiflora*, 341; grandiflora, 342; Halei, 265; *Hanseni*, 257; hastata, 278; *hastata* var. *oblongifolia*, 267, var. *paniculata*, 278, f. *rosea*, 278, var. *pinnatifida*, 278; *hirsuta*, 268; *imbricata*, 304; incarnata, 342; *inconspicua*, 338; integrifolia, 342; *Lamberti*, 316, var. *rosea*, 316; *lasiostachys*, 290; *leucanthemifolia*, 265; lilacina, 340; litoralis, 257; *litoralis* var. *brasiliensis*, 255, var. *caracasana*, 257; *littoralis β. leptostachya*, 257, *α. pycnostachya*, 257; *longiflora*, 315; longifolia, 272; *Lucaeana*, 293; MacDougalii, 288; *MacDougalii* mut. *rosella*, 288; macrodonta, 289, *358, 360;* maritima, 320; Matthesii, 342; menthaefolia, 263; *mollis*, 268, 286; *moranensis*, 318; neomexicana, 296, var. hirtella, 298, var. xylopoda, 297; *Oblaetia*, 315; *Obletia*, 315; officinalis, 262; *officinalis* var. *hirsuta*, 296; Orcuttiana, 284; *paniculata*, 278, var. *pinnatifida*, 278; *pauciflora*, 268; *paucifolia*, 268; paucifolia, 342; perennis, 299; *pinnatifida*, 278; plicata, 294; *polystachya*, 268; prostrata, 290; *pu-*

bera, 332; *pulchella*, 332; pumila, 338; *pumila* f. *albiflora*, 313; *quadrangularis*, 254; quadrangulata, 313; racemosa, 332; recta, 271; *remota*, 300; repens, 342; *rigens*, 286; rigida, 252; riparia, 267; robusta, 292; *Roemeriana*, 302; *rubra*, 316; *rudis*, 304; *rugosa*, 282; scabra, 272; *scabra*, 272; scabrella, 343; **setacea**, 339, *362; setosa*, 263; simplex, 282; **sphaerocarpa**, 256; *spuria*, 262; *squarrosa*, 304; stricta, 285; *stricta* f. *albiflora*, 286, β. ? *mollis*, 286, f. *roseiflora*, 286; *strigosa*, 293; subuligera, 303; tampensis, 321; teucriifolia, 334, var. **corollulata**, 335; trifida, 342; **tumidula**, 322; urticifolia, 275; *urticifolia* var. *riparia*, 267, var. *simplex*, 275; *venosa*, 252; *verna*, 337, var. *fissa*, 335; *veronicaefolia*, 268; Wrightii, 328, *358;* xutha, 293

W

Woodsia, 68; obtusa, 68
Woodson, Robert E., Jr. Studies in the Apocynaceae. IV, 605; George H. Pring, and. A new yellow Nymphaea from tropical Africa, 1
Woodwardia, 69; *angustifolia*, 69; areolata, 69
Wynd, F. Lyle. Nutrient solutions for orchids, 363; The sensitivity of orchid seedlings to nutritional ions, 223; Sources of carbohydrate for germination and growth of orchid seedlings, 569

X

Xanthocephalum, 24; linearifolium, 25
Xanthoria polycarpa, 15
Xylose, growth of orchid seedlings in solutions containing, 574, 576

Y

Yeast-like organisms, 471; cytology of, 477

Z

Zapania bracteosa, 304
Zygadenus elegans, 801